

Associations Between Coinfection Prevalence of *Borrelia lusitaniae*, *Anaplasma* sp., and *Rickettsia* sp. in Hard Ticks Feeding on Reptile Hosts

Radovan Václav · Martina Ficová · Pavol Prokop ·
Tatiana Betáková

Received: 17 May 2010 / Accepted: 3 August 2010 / Published online: 14 August 2010
© Springer Science+Business Media, LLC 2010

Abstract An increasing number of studies reveal that ticks and their hosts are infected with multiple pathogens, suggesting that coinfection might be frequent for both vectors and wild reservoir hosts. Whereas the examination of associations between coinfecting pathogen agents in natural host–vector–pathogen systems is a prerequisite for a better understanding of disease maintenance and transmission, the associations between pathogens within vectors or hosts are seldom explicitly examined. We examined the prevalence of pathogen agents and the patterns of associations between them under natural conditions, using a previously unexamined host–vector–pathogen system—green lizards *Lacerta viridis*, hard ticks *Ixodes ricinus*, and *Borrelia*, *Anaplasma*, and *Rickettsia* pathogens. We found that immature ticks infesting a temperate lizard species in Central Europe were infected with multiple pathogens. Considering *I. ricinus* nymphs and larvae, the prevalence of *Anaplasma*, *Borrelia*, and *Rickettsia* was 13.1% and 8.7%, 12.8% and 1.3%, and 4.5% and 2.7%, respectively. The patterns of pathogen prevalence and observed coinfection rates suggest that the risk of tick infection with one pathogen is not independent of other pathogens. Our results indicate that *Anaplasma* can play a role in suppressing the transmission of *Borrelia* to tick vectors. Overall, however, positive effects of

Borrelia on *Anaplasma* seem to prevail as judged by higher-than-expected *Borrelia*–*Anaplasma* coinfection rates.

Introduction

An increasing number of studies report that ticks and their vertebrate hosts are infected with a mixture of pathogens, suggesting that coinfection might be the rule rather than the exception for both vectors and wild reservoir hosts [5, 14, 45]. Because direct or indirect relationships within a community of pathogens inevitably affect their fitness, coinfection is expected to result in synergic or antagonistic effects between the members of a pathogen community [26]. Indeed, some pathogens can evade the host's immune system more easily if they are transmitted by vectors simultaneously with other infections [52]. Hosts coinfecting with different pathogens may show more severe symptoms of diseases [22], and host coinfection status can modulate the transmission of pathogens to vectors [47].

Ginsberg [14] has shown that if tick populations are large and individual hosts are in their life infested by multiple ticks, the probability of host exposure to a pathogen is close to 100% even if the proportion of infected vectors is small. While the probability of infection could further increase with vector coinfection, this might not be the case if there are negative interactions between coinfecting pathogens [47]. The order of infection also can affect the risk of infection because some pathogens can be more immunosuppressive than others and thus modulate the probability of consecutive infections [19]. Therefore, it is important whether coinfection occurs simultaneously (true coinfection) or sequentially because the two scenarios can

R. Václav (✉) · P. Prokop
Institute of Zoology, Slovak Academy of Sciences,
Dubravská cesta 9,
84506 Bratislava, Slovakia
e-mail: Radovan.Vaclav@savba.sk

M. Ficová · T. Betáková
Virological Institute, Slovak Academy of Sciences,
Dubravská cesta 9,
84505 Bratislava, Slovakia

have different clinical and epidemiological outcomes [5, 26]. Whereas the examination of associations between coinfecting pathogens in natural host–vector–pathogen systems is a prerequisite for better insight into disease ecology, only a few studies have explicitly examined the associations between pathogens in wild vector or host populations [11, 32].

Reptiles represent an excellent group to study the epidemiology of various pathogens because of their long life span, site faithfulness, seasonal nature of immunity, and distinct modes of reproduction [43] and immune responses [23]. Moreover, lizards, including green lizards *Lacerta viridis*, are locally important hosts for the early life stages of hard ticks [7, 49]. Though lizards have been long considered as non-competent hosts for *Borrelia* spirochetes, recent experimental and empiric work strongly suggests that lizards can be competent reservoirs of a potentially pathogenic *Borrelia* genospecies, *Borrelia lusitaniae* [2, 9, 12, 29, 42]. In contrast to *Borrelia* spirochetes, little is known about the prevalence and ecology of other tick-borne pathogens associated with reptile hosts. We examined the patterns of associations between three different pathogen agents under natural conditions, using a previously unexamined host–vector–pathogen system. Specifically, we used the green lizard *L. viridis*, which serves as a vertebrate host for hard ticks, the hematophagous ectoparasites—known vectors of several important pathogens, as well as the three selected pathogens: *Borrelia burgdorferi* sensu lato, *Anaplasma* sp., and *Rickettsia* sp.

Materials and Methods

Study System and Area

The European green lizard is a 35–40-cm (total length) long lizard, inhabiting warm and dry areas of central Europe and the Balkans. The breeding season of green lizards begins after hibernation in April and lasts until early June when females lay about seven to 20 eggs. The lizard is one of the largest European lacertids and has a relatively long lifespan (>9 years). *Ixodes ricinus* (Acari: Ixodidae) is an opportunistic hematophagous ectoparasite species of hard ticks, commonly infesting reptiles [4]. *Borrelia* spirochetes are bacteria transmitted by ticks and causing inflammatory disorders [36]. The members of *Anaplasmataceae* are tick-borne obligatory intracellular bacterial parasites attacking neutrophils [13]. Similar to *Anaplasma*, *Rickettsiae* are obligatory intracellular bacterial parasites, but they are transmitted by different types of hematophagous arthropods (ticks, mites, lice, and fleas), which can, due to transovarian transmission of bacteria, serve as both vectors and hosts

[31]. The study site (48°57' N, 20°44' E, ~200 to 400 m above sea level) in the Slovak Karst National Park was ~10 ha large, characterized by warm temperate climate, open habitat and broad-leaved scrub vegetation (*Quercus petraea*, *Carpinus betulus*, *Cornus mas*, *Acer campestre*, *Crataegus monogyna*, and *Prunus mahaleb*).

Sample Collection

In 2007, we captured free-living green lizards by slipnoosing between 6 April and 11 May. During four 3–4-day-long trips, we trapped a total of 89 individuals. Each captured lizard was sexed according to throat coloration, head shape, the size of femoral pores, the shape of the ventral side of tail base (male tail base has two bulges because it contains the paired hemipenes), and bite marks on the head and belly (males bite the belly of females during copulation; head bite marks usually come from territorial disputes between males). All lizards were marked with a marker pen in the field and stored in individual containers. Later in the same day, we removed with forceps all the ticks feeding on lizards. In addition, we obtained a tissue sample by clipping the tip of one toe on the rear leg from each lizard. Both ticks and toe tips were stored in 93% ethanol. A drop of blood obtained following toe clipping was used for blood smears (not analyzed here). All lizards were released exactly at the point of capture the morning after their capture. The life stage and species of collected ticks was determined using microscope and standard identification keys [8].

DNA Extraction and Detection of *Borrelia*, *Rickettsia*, and *Anaplasma* by PCR

DNA extraction from samples was based on the method using an anion chelating resin (Chelex 100® Resin, Bio-Rad Laboratories, CA, USA, [38]). Briefly, tick was transferred into a 1.5-ml microcentrifuge tube containing 100 µl of pure water, 0.005 g Chelex 100 resin, and 6 µl proteinase K (10 µg/ml). The mixture was homogenized and vortexed 2 min at room temperature. The homogenized sample was then centrifuged at 13,000×g for 1 min and incubated at 96°C for 35 min. After vortexing and centrifugation at 13,000×g for 1 min, the sample was incubated at 96°C for 15 min. The mixture was then centrifuged at 13,000×g for 3 min, and the supernatant was used as a template in PCR essays.

2xPCR Master Mix (Fermentas) was used for all PCR reactions. Primers specific for all *Borrelia* species, together with PCR conditions, have been described previously by Kurtenbach et al. [24]. Rickettsial DNA was detected by PCR as described previously by using primers Rp877p and Rp1258r, which amplify a 396-bp fragment of the citrate

synthase gene (*gltA*) of *Rickettsia* [44]. The DNA extracted from the ticks was also screened with primers EHR16SR and EHR16SD, which amplify a 345-bp fragment of the 16S rRNA gene of bacteria within the family Anaplasmataceae, including the genera *Anaplasma*, *Ehrlichia*, *Neorickettsia*, and *Wolbachia* [37]. All specific PCR products were separated by electrophoresis in 2% agarose gels and visualized by staining with ethidium bromide. *Anaplasma* and *Rickettsia* PCR products were sequenced for further determination using a commercial sequencing service (BITCET).

Reverse Line Blot Hybridization

The biotinylated spacer fragment of the *Borrelia* PCR was hybridized with six *B. burgdorferi* genospecies-specific oligonucleotide probes: *B. burgdorferi* sensu lato, *B. burgdorferi* sensu stricto, *Borrelia valaisiana*, *Borrelia garinii*, *Borrelia afzelii*, and *B. lusitaniae*. Biotinylated *Rickettsia* PCR products were examined using the genospecies-specific oligonucleotide probes to *Rickettsia slovacica* and *Rickettsia helvetica* as described by Jado et al. [21].

The reverse line blotting primers and technique has been in detail described elsewhere [30, 50]. Briefly, solutions with 100 pmol of five different amino-linked oligonucleotide probes were coupled covalently to an activated Bodine C membrane in a line pattern by using a miniblotter. After binding of the oligonucleotide probes, the membrane was taken from the miniblotter, washed in 2× SSPE (360 mM NaCl, 20 mM Na₂HPO₄×H₂O, 2 mM EDTA) with 0.1% sodium dodecyl sulfate (SDS) at RT, and again placed in the miniblotter with the oligonucleotide lines perpendicular to the slots. Ten microliters of the biotin-labeled PCR product was diluted in 150 ml of 2× SSPE–0.1% SDS, denatured for 10 min at 99°C, and cooled rapidly on ice. The slots of the miniblotter were filled with the denatured PCR product, and hybridization was performed for 1 h at 45°C. The membrane was removed from the miniblotter and was washed twice for 10 min each time in 2× SSPE–0.1% SDS at 40°C. Subsequently, the membrane was incubated for 30 min at 40°C with streptavidin-horse radish peroxidase (Novagen) diluted 1:4,000 in 2× SSPE–0.5% SDS and was washed twice for 10 min in 2× SSPE–0.5% SDS. Hybridization was visualized by labeling with TMB Stabilized Substrate for Horseradish Peroxidase (Promega).

PCR and Restriction Fragment Length Polymorphism Analysis

All samples testing positive for *Borrelia* were further characterized by 5S–23S rRNA integric spacer restriction fragment length polymorphism (RFLP) analysis. Primers

and conditions have been described elsewhere [20, 39]. PCR products were digested with *Mse*I and *Dra*I, according to manufacturer instructions (Fermentas, Merck). Digested DNA was electrophoresed through 16% polyacrylamide gels and stained with ethidium bromide; 100 bp DNA ladder (Promega) was used as a molecular size marker.

Data Analyses

Generalized linear mixed models (GLMM) were used to examine variation in the occurrence of pathogens, where lizard identity, and the interaction of lizard identity and the capture date were used as random factors. The residual random effect was included in models as an overdispersion parameter [28]. Parameter estimates and 95% confidence limits were calculated with SAS (SAS 2002 Institute, Cary, NC, USA) using the PROC GLIMMIX procedure. Binomial distribution and the logit link function were applied. The Satterthwaite method, which can give non-integer values, was used to calculate degrees of freedom [28]. The significance of fixed effects was based on the *F* statistics. The interaction terms involving relevant pathogen pairs (i.e., *Borrelia* × *Rickettsia*, *Borrelia* × *Anaplasma*, or *Anaplasma* × *Rickettsia*, respectively) were included in GLMM. Concerning the removal of interaction terms from models and the correct interpretation of main effects, we followed the recommendations of Brambor et al. [6]. Briefly, instead of showing the statistics for the main effects as average (unconditional), to assess the effect of constitutive terms in models including interactions, we show and interpret their effects as conditional [40]. The *G* test was used to test the differences between observed and expected coinfection rates. Expected coinfection rates were calculated as the product of the prevalence of each pathogen individually [32].

Results

PCR Detection of *Borrelia*, *Anaplasma*, and *Rickettsia* DNAs in Ticks and Lizards

Various PCRs were applied to DNA extracts from 799 *Ixodes ricinus* and 25 *Dermacentor marginatus* ticks collected from 89 green lizards. Adult ticks were not found on the lizards. The most abundant pathogen detected in the ticks was *Anaplasma* sp. (Table 1). The further sequencing of *Anaplasma* sp. PCR products revealed 93–96% homology with uncultured *Anaplasma* sp. clone HLA344 16S ribosomal RNA gene (GU075704).

The prevalence of *Borrelia* in *I. ricinus* was only slightly lower compared with *Anaplasma* sp. (Table 1). All

Table 1 Prevalence of *Anaplasma* sp., *Borrelia lusitaniae*, and *Rickettsia* sp. in immature *Ixodes ricinus* and *Dermacentor marginatus* ticks feeding on green lizards *Lacerta viridis*

Tick species	<i>Anaplasma</i> sp.		<i>Borrelia lusitaniae</i>		<i>Rickettsia</i> sp.		No. nymphs/ larvae
	No. (%) positive nymphs	No. (%) positive larvae	No. (%) positive nymphs	No. (%) positive larvae	No. (%) positive nymphs	No. (%) positive larvae	
<i>Ixodes ricinus</i>	85 (13.1%)	13 (8.7%)	83 (12.8%)	2 (1.3%)	29 (4.5%)	4 (2.7%)	649/150
<i>Dermacentor marginatus</i>	1 (7%)	1 (10%)	1 (7%)	0 (0%)	1 (7%)	0 (0%)	15/10

Adult ticks of neither species were found on green lizards

Borrelia-positive samples were subjected to further genotyping analyses.

DNA of *Rickettsia* was detected in 33 ticks. Sequence analysis of *Rickettsia* sp. PCR products identified 99–100% homology with *Rickettsia* sp. 362 23S ribosomal gene (DQ139797.1) and 93% homology with *Rickettsia massiliae* MTU5 (CP000683.1). One *Rickettsia* sp. PCR product was determined as *R. helvetica* by reverse line blot (RLB).

Seventeen nymphs contained both *Borrelia* and *Anaplasma*, ten nymphs contained *Borrelia* and *Rickettsia*, and four nymphs contained *Anaplasma* and *Rickettsia*. We have also observed triple infection with all three pathogens for four nymphs (Fig. 1).

DNA extracted from 25 *D. marginatus* ticks tested positive for *Anaplasma* sp., *B. burgdorferi* s.l., and *Rickettsia* sp. (Table 1).

As for the host, 47.19% (42/89), 51.69% (46/89), and 22.47% (20/89) of different green lizards carried at least one feeding tick testing positive for *B. lusitaniae*, *Anaplasma* sp., or *Rickettsia* sp., respectively. However, examining the toe tips of 89 lizards, we detected neither *B. burgdorferi* s.l., nor *Anaplasma* sp. and *Rickettsia* sp. in any lizard toe tissue sample.

Identification of *Borrelia* Using RLB and RFLP Analyses

RLB was designed to differentiate *Borrelia* genospecies. To assess the specificity of the assay, control samples were used in the RLB assay under stringent hybridization conditions. All our samples tested were positive for *B. burgdorferi* s.l. and *B. lusitaniae*. Cross-hybridization between different genospecies did not occur.

All samples were also analyzed by PCR-RFLP methods of rrf(5S)–rrl(23S) rRNA *B. burgdorferi* s.l. PCR amplicons. RFLP analysis of the amplified products yielded two different patterns. The most abundant was the pattern similar to that of *B. lusitaniae* PotiBI, pattern E [10]. Only three investigated samples showed a slightly different pattern with one extra restriction site. The similar pattern was described for the sample GT163, pattern

F + E, determined also as *B. lusitaniae* [10]. Overall, all detected *Borreliae* belonged to the genospecies *B. lusitaniae*.

Tick-Vector Infection Patterns

The rates of *I. ricinus* single infection versus coinfection differed significantly between the three types of pathogens




Figure 1 Schematic illustration of *I. ricinus* infection patterns. The values correspond to the number of *I. ricinus* ticks infected with different combinations of three pathogens: *B. lusitaniae*, *Anaplasma* sp., and *Rickettsia* sp. Shown are infection patterns in **a** 649 nymphs and **b** 150 larvae

($G_2=10.34$, $p<0.05$). This was due to *Rickettsia* (Rc), which, compared with other pathogens (i.e., *B. lusitaniae* (Bl) and *Anaplasma* sp. (An)), occurred in ticks as coinfection more often than single infection (Fig. 1). In addition, the nymphal *I. ricinus* coinfection rates of (1) Bl with An, (2) Bl with An and Rc, and (3) Bl with Rc were significantly higher than expected based on single tick infection rates (Table 2).

Patterns of Within-Tick Infection in *I. ricinus*

The probability of Bl infection did not differ between dates and host sexes (GLMM, date: $F_{3,81.44}=0.70$, $p>0.05$; sex: $F_{1,132.9}=0.56$, $p>0.05$), but it was significantly higher for tick nymphs than larvae ($F_{1,747.7}=14.62$, $p<0.05$). The effect on the occurrence of Bl infection of An tick infection was conditional on Rc, but not the other way around. Namely, if ticks were infected with Rc, the likelihood of Bl occurrence was higher regardless of An coinfection status (An × Rc: $F_{1,785.5}=1.21$, $p>0.05$; effect of Rc occurrence sliced by An occurrence, An absent: $F_{1,735}=30.87$, $p<0.05$; An present: $F_{1,783.8}=5.03$, $p<0.05$; Fig. 2). In contrast, if ticks were infected with An, the likelihood of Bl tick infection was lower when An was infecting ticks alone compared with cases when An-infected ticks were coinfecting with Rc (effect of An occurrence sliced by Rc occurrence, Rc absent: $F_{1,788.8}=14.12$, $p<0.05$; Rc present: $F_{1,782.3}=0.06$, $p>0.05$; Fig. 2).

Trapping date, host sex, and tick life stage did not significantly predict the occurrence of An tick infection (GLMM, date: $F_{3,107.6}=1.75$, $p>0.05$; host sex: $F_{1,92.82}=0.49$, $p>0.05$; tick stage: $F_{1,735.8}=0.99$, $p>0.05$). The

Table 2 Observed and expected coinfection prevalence of *Anaplasma* sp. (An), *Borrelia lusitaniae* (Bl), and *Rickettsia* sp. (Rc) for *Ixodes ricinus* ticks feeding on green lizards *Lacerta viridis*

Coinfection type	Samples	Observed coinfecting (%)	Expected coinfecting (%)	G	p
An × Bl					
Larvae	150	0.00	0.05	0.00	1.000
Nymphs	649	2.62	0.74	6.15	0.008 ^a
Bl × Rc					
Larvae	150	0.00	0.01	0.00	1.000
Nymphs	649	1.54	0.14	7.41	0.003 ^a
An × Rc					
Larvae	150	0.67	0.16	1.39	0.238
Nymphs	649	0.62	0.15	1.93	0.164
Bl × An × Rc					
Larvae	150	0.00	0.00	0.00	1.000
Nymphs	649	0.62	0.01	5.56	0.018 ^a

^a Statistically significant at $p\leq 0.05$




Figure 2 Probability of *B. lusitaniae* infection in *I. ricinus* as a function of *Anaplasma* and *Rickettsia* coinfection prevalence. Displayed are the estimates of least-squares means and 95% confidence limits obtained by GLMM

likelihood of An tick infection was significantly higher if ticks were infected with Bl, Rc, or both pathogens compared with cases when ticks were not infected with the two pathogens (Bl × Rc: $F_{1,767.4}=1.54$, $p>0.05$; effect of Bl occurrence sliced by Rc occurrence, Rc absent: $F_{1,714.6}=13.37$, $p<0.05$, Rc present: $F_{1,772.3}=0.02$, $p>0.05$; effect of Rc occurrence sliced by Bl occurrence, Bl absent: $F_{1,772.1}=5.95$, $p<0.05$, Rc present: $F_{1,755.5}=0.18$, $p>0.05$; Fig. 3).

The probability of Rc tick infection did not differ between host sexes, trapping date, and tick life stage (GLMM, date: $F_{3,134.8}=2.09$, $p>0.05$; host sex: $F_{1,136.5}=0.02$, $p>0.05$; tick stage: $F_{1,774.9}=0.26$, $p>0.05$). Considering An infection status, the probability of Rc infection was analogous to that




Figure 3 Probability of *Anaplasma* infection in *I. ricinus* as a function of *Borrelia* and *Rickettsia* coinfection prevalence. Displayed are the estimates of least-squares means and 95% confidence limits obtained by GLMM

of BI infection. Particularly, if ticks were infected with An, the likelihood of Rc tick infection was lower if An was infecting ticks alone compared with cases when An-ticks were coinfecting with BI (BI \times An: $F_{1,788.4}=7.08$, $p<0.05$; effect of An occurrence sliced by BI occurrence, BI absent: $F_{1,787.5}=14.77$, $p<0.05$; BI present: $F_{1,790}=0.21$, $p>0.05$; Fig. 4). In turn, if BI was detected in ticks, the probability of Rc infection was higher regardless of An infection status (effect of BI occurrence sliced by An occurrence, An absent: $F_{1,776.4}=62.91$, $p<0.05$; An present: $F_{1,789.3}=4.37$, $p<0.05$; Fig. 4).

Discussion

To our knowledge, this is the first study reporting *Anaplasma* in ticks feeding on a scaled reptile (Reptilia: Squamata) from Central Europe. *Anaplasma* was recently detected in ticks infesting western fence lizards (*Sceloporus occidentalis*) and northern alligator lizards (*Elgaria coerulea*) from North America [33], sand lizards *Lacerta agilis* from West Europe [48], and savannah monitors *Varanus exanthematicus* from West Africa [34]. In addition, this work is one of the few studies reporting *Rickettsia* in ticks from reptile hosts [15, 41, 51]. Our study is first to show not only that hard ticks feeding on green lizards are infected with both *Borrelia* and *Anaplasma* (see [32] for other host–vector systems) but also that there are distinct patterns of pathogen coinfection prevalence in *I. ricinus* collected from lizards.

Our work has revealed that ticks feeding on green lizards are not infrequently infected with *Borrelia* spirochetes, *Anaplasma* sp., and *Rickettsia* sp. (Table 1).




Figure 4 Probability of *Rickettsia* infection in *I. ricinus* as a function of *Borrelia* and *Anaplasma* tick coinfection prevalence. Displayed are the estimates of least-squares means and 95% confidence limits obtained by GLMM

Our study corroborates the recent results on the strong association of *B. lusitaniae* genospecies with lizard hosts [2, 12, 29, 42]. Similar to the results by Richter and Matuschka [42], who examined *B. lusitaniae* infection of *I. ricinus* ticks infesting *L. agilis*, we did not detect any other *Borrelia* genospecies than *B. lusitaniae*. Yet, using RFLP analysis, we detected in ticks two patterns of *B. lusitaniae*—pattern E and pattern F + E.

Our result on 47% of lizards being infested with at least one tick infected with *B. lusitaniae* lies between 66% reported by Richter and Matuschka [42] and 29% reported by Majláthová et al. [29]. The discrepancy between this and the latter study, though both being conducted on the same lizard species and in the same geographic region, could be due to the differences in the period of tick collection. While Majláthová et al. [29] collected ticks from lizards mostly outside the lizards' breeding season (May–September), our study was conducted in the spring during the period of the most intensive sexual activity of lizards (April–May), when the immune system is compromised by sex hormones and lower ambient temperature and lymphoid organs are not yet well developed after winter hibernation [35].

The nature of our study makes it impossible to confirm that the infected ticks acquired *Borrelia* or other pathogens from lizards on which they fed. We collected the tip of one toe from each lizard for molecular analysis. However, we did not detect *Borrelia* in any of these samples. This result adds to the negative results on the use of keratinized tissue parts (tail tips) of lizards [2, 29]. Nevertheless, the fact that no tick was infected with other *Borrelia* genospecies than *B. lusitaniae*, even though in Slovakia the majority of questing ticks and the ticks feeding on rodent or avian hosts are infected with other *Borrelia* genospecies [16–18, 25, 29] (Table 3), strongly suggests that the examined ticks obtained *Borrelia* from lizards. Further work needs to focus on other tissues and serological analysis to detect *Borrelia* in lizard hosts and validate that *Borrelia* can be transmitted from green lizards to ticks.

Despite the wide confidence limits and relatively few ticks coinfecting with multiple pathogens, our study has revealed that the prevalence of both *B. lusitaniae* and *Rickettsia* in ticks feeding on green lizards was significantly lower if the ticks infected with *Anaplasma* were not coinfecting with *Rickettsia* or *B. lusitaniae*, respectively. In contrast, the probability of *Anaplasma* tick infection increased if the ticks were infected with *B. lusitaniae* and/or *Rickettsia*. Therefore, it appears that there is an important negative effect (direct or indirect) of *Anaplasma* on *B. lusitaniae* and *Rickettsia* prevalence, a positive impact of *B. lusitaniae* and *Rickettsia* on *Anaplasma* prevalence, and positive interactions between *B. lusitaniae* and *Rickettsia*.

Table 3 Patterns of host infestation by ticks and tick infection for different vertebrate hosts in the study region

Vertebrate host	Tick vector	Hosts infested by ticks (%) ^a	Average tick load	Hosts infested by infected ticks (%) ^b	Infected ticks (%)	Detected pathogens infecting ticks	Reference
Lizards							
<i>Lacerta viridis</i>	<i>Ixodes ricinus</i>	88	2.8	29	16.6	<i>Borrelia lusitaniae</i> , <i>B. afzelii</i> , <i>B. garinii</i> , <i>B. burgdorferi</i> s.s., <i>B. valaisiana</i>	[29]
<i>Lacerta viridis</i>	<i>Ixodes ricinus</i>	93	8.3	71.9	22.0	<i>B. lusitaniae</i> , <i>Anaplasma</i> sp., <i>Rickettsia</i> sp.	This study
<i>Lacerta viridis</i>	<i>Dermacentor marginatus</i>	13	0.3	4.2	16.0	<i>B. burgdorferi</i> sensu lato, <i>Anaplasma</i> sp., <i>Rickettsia</i> sp.	This study
Birds							
<i>Turdus merula</i>	<i>Ixodes ricinus</i>	84	17.4	50/76.9	35.9	<i>B. garinii</i> , <i>B. valaisiana</i> , <i>B. afzelii</i> , <i>B. burgdorferi</i> sensu stricto	[46]
<i>Turdus philomelos</i>	<i>Ixodes ricinus</i>	81	11.7	50/63.6	38.1	<i>B. garinii</i> , <i>B. valaisiana</i> , <i>B. afzelii</i>	[46]
Mammals ^c							
<i>Apodemus flavicollis</i>	<i>Ixodes ricinus</i>	73/14	5.2	24.6/46.2	12.7	<i>B. afzelii</i> , <i>B. valaisiana</i> , <i>B. burgdorferi</i> s.s.	[17]
<i>Clethrionomys glareolus</i>	<i>Ixodes ricinus</i>	62/11	2.5	27.9/41.7	30.3	<i>B. afzelii</i>	[17]

^a Percentages of infested mammal hosts refer to hosts infested by larvae/nymphs

^b Percentages of bird and mammal hosts infested by infected ticks refer to hosts infested by larvae/nymphs

^c Results for mammals are from Western Slovakia

Only a few studies report negative relationships between different vector-borne pathogens [1, 30], and most of these works report the associations between different strains of the same pathogen such as *Anaplasma* [14]. There is currently no convincing evidence that the interactions between *B. burgdorferi* and *Anaplasma phagocytophilum* may take place within vectors [26, 27]. In contrast, Levin [26] has demonstrated that such interactions, likely brought about by host immune responses, can occur within the host. The author found that if white-footed mice *Peromyscus leucopus* were primarily infected with *Anaplasma* and then secondarily with *Borrelia*, fewer xenodiagnostic *Ixodes scapularis* ticks acquired infection with *Borrelia* compared with the ticks feeding on mice infected with *Borrelia* only. An analogous result for *Anaplasma* tick infection was obtained if mice were first infected with *Borrelia* and then with *Anaplasma*. The infection pattern of tick larvae (a measure of host spirochete infection, e.g., [3]), with larvae being predominantly infected with *Anaplasma* only, suggests that the negative associations between *Anaplasma* and *Borrelia* and *Rickettsia* tick prevalence resulted from their interactions within tick hosts rather than ticks. Taken together, our work accords with the suggestion by Levin [26] in that *Anaplasma* infection may suppress the development of *Borrelia* spirochetes and suggests that *B. lusitaniae*, and also *Rickettsia*, do not benefit from coinfection with *Anaplasma*.

Existing body of work does not clearly suggest that the rates of observed tick coinfections with *Borrelia* and

Anaplasma are higher than expected [32]. Our study showed that tick *Borrelia*–*Anaplasma*, and also *Borrelia*–*Rickettsia* and *Borrelia*–*Anaplasma*–*Rickettsia* coinfection rates, is higher than expected based on single infection rates. Thus, while within-tick infection patterns have revealed a negative impact of *Anaplasma* on *Borrelia* prevalence, the results on observed coinfection rates suggest that the positive effects of *Borrelia* on *Anaplasma* prevalence in ticks prevail in our study system. In addition, our results on the negative associations between *Anaplasma* and *Borrelia* prevalence, as well as the heterogeneity of results on vector *Borrelia*–*Anaplasma* coinfection rates [32], lend support to the idea that in nature, a substantial number of *Borrelia*–*Anaplasma* coinfections result from interactions between different pathogens and/or sequential (transstadial) rather than simultaneous vector infection [26, 50].

To conclude, we show that ticks infesting a free-living temperate lizard species in a Central European region were infected with multiple pathogens with some of them previously found only in other vertebrates. The patterns of pathogen prevalence and observed coinfection rates suggest that the risk of tick infection with one pathogen is not independent of other pathogens. Overall, positive effects of *Borrelia* on *Anaplasma* seem to prevail as judged by higher-than-expected *Borrelia*–*Anaplasma* coinfection rates. Further laboratory studies are necessary to prove if *Anaplasma* can play a role in suppressing the transmission of *Borrelia* from the host to tick vectors and thus mediate its circulation in nature.

Acknowledgments We thank Dr. Ján Krištofik for tick determination. Milan Olekšák and the directive of the Slovak Karst National Park provided logistic support. This study was funded by the VEGA grants no. 2/7080/27 and 1/0207/08. The study was conducted under approval from the Ministry of Environment of the Slovak Republic (license no. 1430/467/04-5.1).

References

- Alekseev AN, Burenkova LA, Vasil'eva IS, Dubinina EV, Chunikhin SP (1996) The functioning of foci of mixed tick-borne infections on Russian territory. *Med Parazitol* 4:9–16 (In Russian)
- Amore G, Tomassone L, Grego E, Ragagli C, Bertolotti L, Nebbia P, Rosati S, Mannelli A (2007) *Borrelia lusitaniae* in immature *Ixodes ricinus* (Acari: Ixodidae) feeding on common wall lizards in Tuscany, Central Italy. *J Med Entomol* 44:303–307
- Barbour A (2004) Specificity of *Borrelia*-tick vector relationships. In: Gillespie SH, Smith GL, Osbourn A (eds) *Microbe–vector interactions in vector-borne diseases*. Cambridge University Press, Cambridge, pp 75–90
- Barnard SM, Durden LA (2000) A veterinary guide to the parasites of reptiles, vol. 2. Arthropods (excluding mites). Krieger, Malabar
- Belongia EA (2002) Epidemiology and impact of coinfections acquired from *Ixodes* ticks. *Vector-Borne Zoonot Dis* 2:265–273
- Brambor T, Clark WR, Golder M (2005) Understanding interaction models: improving empirical analyses. *Polit Anal* 13:1–20
- Casher LE, Lane RS, Barrett RH, Eisen L (2002) Relative importance of lizards and mammals as hosts for ixodid ticks in northern California. *Exp Appl Acarol* 26:127–143
- Daniel M, Černý V (1971) *Klíč zvířeny ČSSR*. Československá Akademie Vied, Praha
- De Carvalho IL, Fonseca JE, Marques JG, Ullmann A, Hojgaard A, Zeidner N, Nuncio MS (2008) Vasculitis-like syndrome associated with *Borrelia lusitaniae* infection. *Clin Rheumatol* 27:1587–1591
- De Michelis S, Sewell HS, Collares-Pereira M, Santos-Reis M, Schouls LM, Benes V, Holmes EC, Kurtenbach K (2000) Genetic diversity of *Borrelia burgdorferi* sensu lato in ticks from mainland Portugal. *J Clin Microbiol* 38:2128–2133
- Dib L, Bitam I, Tahri M, Bensouilah M, De Meeüs T (2008) Competitive exclusion between Piroplasmosis and Anaplasmosis agents within Cattle. *PLoS Pathog* 4:e7
- Dsouli N, Younsi-Kabachii H, Postic D, Nouira S, Gern L, Bouattour A (2006) Reservoir role of lizard *Psammotromus algirus* in transmission cycle of *Borrelia burgdorferi* sensu lato (Spirochaetaceae) in Tunisia. *J Med Entomol* 43:737–742
- Dumler JS, Choi K-S, Garcia-Garcia JC, Barat NS, Scorpio DG, Garyu JW, Grab DJ, Bakken JS (2005) Human granulocytic Anaplasmosis and *Anaplasma phagocytophilum*. *Emerg Infect Dis* 11:1828–1834
- Ginsberg HS (2008) Potential effects of mixed infections in ticks on transmission dynamics of pathogens: comparative analysis of published records. *Exp Appl Acarol* 46:29–41
- Graves S, Stenos J (2003) *Rickettsia honei*: a spotted fever group rickettsia on three continents. *Ann New York Acad Sci* 990:62–66
- Gronesova P, Ficova M, Mizakova A, Kabat P, Trnka A, Betakova T (2008) Prevalence of avian influenza viruses, *Borrelia garinii*, *Mycobacterium avium*, and *Mycobacterium avium* subsp. *paratuberculosis* in waterfowl and terrestrial birds in Slovakia, 2006. *Avian Pathol* 37:537–543
- Hanincová K, Schäfer SM, Etti S, Sewell H-S, Taragelová V, Ziak D, Labuda M, Kurtenbach K (2003) Association of *Borrelia afzelii* with rodents in Europe. *Parasitol* 126:11–20
- Hanincová K, Taragelová V, Koci J, Schäfer SM, Hails R, Ullmann AJ, Piesman J, Labuda M, Kurtenbach K (2003) Association of *Borrelia garinii* and *B. valaisiana* with songbirds in Slovakia. *Appl Environ Microbiol* 69:2825–2830
- Holden K, Hodzic E, Feng S, Freet KJ, Lefebvre RB, Barthold SW (2005) Coinfection with *Anaplasma phagocytophilum* alters *Borrelia burgdorferi* population distribution in C3H/HeN mice. *Infect Immun* 73:3440–3444
- Ishiguro F, Takada N, Masuzawa T, Fukui T (2000) Prevalence of Lyme disease *Borrelia* spp. in ticks from migratory birds on the Japanese mainland. *Appl Environ Microbiol* 66:982–986
- Jado I, Escuerdo R, Gil H, Jiménez-Alonso MI, Sousa R, García-Pérey AL, Rodréguéy-Vargas M, Lobo B, Anda P (2006) Molecular method for identification of Rickettsia species in clinical and environmental samples. *J Clin Microbiol* 44:4572–4576
- Krause PJ (2002) Babesiosis. *Med Clin N Am* 86:361–373
- Kuo MM, Lane RS, Giclas PC (2000) A comparative study of mammalian and reptilian alternative pathway of complement-mediated killing of the Lyme disease spirochete (*Borrelia burgdorferi*). *J Parasitol* 86:1223–1228
- Kurtenbach K, Peacey M, Rijpkema SG, Hoodless AN, Nuttall PA, Randolph SE (1998) Differential transmission of the genospecies of *Borrelia burgdorferi* sensu lato by game birds and small rodents in England. *Appl Environ Microbiol* 64:1169–1174
- Kurtenbach K, De Michelis S, Sewell H-S, Etti S, Schäfer M, Hails R, Collares-Pereira M, Hanincová K, Labuda M, Bormane A, Donaghy M (2001) Distinct combinations of *Borrelia burgdorferi* sensu lato genospecies found in individual questing ticks from Europe. *Appl Environ Microbiol* 67:4926–4929
- Levin ML (2007) Effects of coinfection with *Borrelia burgdorferi* and *Anaplasma phagocytophilum* in vector ticks and vertebrate hosts. In: van Nitch P (ed) *Research on Lyme disease*. Nova, New York, pp 1–34
- Levin ML, Fish D (2000) Acquisition of coinfection and simultaneous transmission of *Borrelia burgdorferi* and *Ehrlichia phagocytophila* by *Ixodes scapularis* ticks. *Infect Immun* 68:2183–2186
- Littell RC, Milliken GA, Stroup WW, Wolfinger RD, Schabenberger O (2006) SAS® for mixed models, 2nd edn. SAS Institute Inc, Cary
- Majláthová V, Majláth I, Derdákova M, Vichová B, Pet'ko B (2006) *Borrelia lusitaniae* and green lizards (*Lacerta viridis*), Karst Region, Slovakia. *Emerg Infect Dis* 12:1895–1901
- Mather TN, Piesman J, Spielman A (1987) Absence of spirochetes (*Borrelia burgdorferi*) and piroplasms (*Babesia microti*) in deer ticks (*Ixodes dammini*) parasitized by chalcid wasps (*Hunterellus hookeri*). *Med Vet Entomol* 1:3–8
- Murray PR, Rosenthal KS, Pfaller MA (2009) *Medical microbiology*, 6th edn. Mosby, Philadelphia
- Nieto NC, Foley JE (2009) Meta-analysis of coinfection and coexposure with *Borrelia burgdorferi* and *Anaplasma phagocytophilum* in humans, domestic animals, wildlife, and *Ixodes ricinus*-complex ticks. *Vector-Borne Zoonotic Dis* 9:93–102
- Nieto NC, Foley JE, Bettaso J, Lane RS (2009) Reptile infection with *Anaplasma phagocytophilum*, the causative agent of granulocytic anaplasmosis. *J Parasitol* 95:1165–1170
- Nowak M, Cienuch S, Stańczak J, Siuda K (2010) Detection of *Anaplasma phagocytophilum* in *Amblyomma flavomaculatum* ticks (Acari: Ixodidae) collected from lizard *Varanus exanthematicus* imported to Poland. *Exp Appl Acarol* 51:363–371

35. Origgi F (2007) Reptile immunology. In: Jacobson ER (ed) Infectious diseases and pathology of reptiles. Color atlas and text. CRC, Boca Raton, pp 131–166
36. Pachner A, Steiner I (2007) Lyme neuroborreliosis: infection, immunity, and inflammation. *Lancet Neurol* 6:544–552
37. Parola P, Roux V, Camicas JL, Baradji I, Brouqui P, Raoult D (2000) Detection of Ehrlichiae in African ticks by PCR. *Trans R Soc Trop Med Hyg* 94:707–708
38. Penchenier L, Simo G, Grébaud P, Nkinin S, Laveissière C, Herder S (2000) Diagnosis of human trypanosomiasis, due to *Trypanosoma brucei gambiense* in central Africa, by the polymerase chain reaction. *Trans R Soc Trop Med Hyg* 94:392–394
39. Postic D, Assous MV, Grimont PA, Baranton G (1994) Diversity of *Borrelia burgdorferi* sensu lato evidenced by restriction fragment length polymorphism of rrf (5S)–rrl (23S) intergenic spacer amplicons. *Int J Sys Bacteriol* 44:743–752
40. Quinn GP, Keough MJ (2007) Experimental design and data analysis for biologists. Cambridge University Press, Cambridge
41. Reeves WK, Durden LA, Dasch GA (2006) A spotted fever group *Rickettsia* from an exotic tick species, *Amblyoma exornatum* (Acari: Ixodidae), in a reptile breeding facility in the United States. *J Med Entomol* 43:1099–1101
42. Richter D, Matuschka FR (2006) Perpetuation of the Lyme disease spirochete *Borrelia lusitaniae* by lizards. *Appl Environ Microbiol* 72:4627–4632
43. Rooney AA (2005) Reptiles: the research potential of an overlooked taxon in immunotoxicology. In: Tryphonas H, Fournier M, Blakley BR, Smits J, Brousseau P (eds) Investigative immunotoxicology. CRC Press, Boca Raton, pp 107–128
44. Roux V, Rydkina E, Ereemeeva M, Raoult D (1997) Citrate synthase gene comparison, a new tool for phylogenetic analysis, and its application for the Rickettsiae. *Int J Sys Bacteriol* 47:252–261
45. Swanson SJ, Neitzel D, Reed KD, Belongia EA (2006) Coinfections acquired from *Ixodes* ticks. *Clin Microbiol Rev* 19:708–727
46. Taragel'ová V, Koči J, Hanincová K, Kurtenbach K, Derdáková M, Ogden NH, Literák I, Kocianová E, Labuda M (2008) Blacbirds and song thrushes constitute a key reservoir of *Borrelia garinii*, the causative agent of borreliosis in Central Europe. *Appl Environ Microbiol* 74:1289–1293
47. Thompson C, Spielman A, Krause PJ (2001) Coinfecting deer-associated zoonoses: Lyme disease, babesiosis, and ehrlichiosis. *Clin Infect Dis* 33:676–685
48. Tijssse-Klasen E, Fonville M, Reimerink JHJ, Spitzen A, Sprong H (2010) Role of sand lizards in the ecology of Lyme and other tick-borne diseases in the Netherlands. *Parasit Vectors* 3:42
49. Václav R, Prokop P, Fekiač V (2007) Expression of breeding coloration in European Green Lizards (*Lacerta viridis*): variation with morphology and tick infestation. *Can J Zool* 85:1199–1206
50. Wielinga PR, Gaasenbeek C, Fonville M, de Boer A, de Vries A, Dimmers W, Akkerhuis OP, Jagers G, Schouls LM, Borgsteede F, van der Giessen JW (2006) Longitudinal analysis of tick densities and *Borrelia*, *Anaplasma*, and *Ehrlichia* infections of *Ixodes ricinus* ticks in different habitat areas in The Netherlands. *Appl Environ Microbiol* 72:7594–601
51. Whitworth T, Popov V, Han V, Bouyer D, Stenos J, Graves S, Ndip L, Walker D (2003) Ultrastructural and genetic evidence of a reptilian tick, *Aponomma hydrosauri*, as a host of *Rickettsia honei* in Australia—possible transovarial transmission. *Ann N Y Acad Sci* 990:67–74
52. Zeidner NS, Dolan MC, Massung R, Piesman J, Fish D (2000) Coinfection with *Borrelia burgdorferi* and the agent of human granulocytic ehrlichiosis suppresses IL-2 and IFN γ production and promotes an IL-4 response in C3H/HeJ mice. *Parasite Immunol* 22:581–588